

There are No Causality Problems for Fermi's Two Atom System

Detlev Buchholz^a and Jakob Yngvason^b

^a II. Institut für Theoretische Physik, Universität Hamburg
Luruper Chaussee 149, D-22761 Hamburg, Germany

^b Science Institute, University of Iceland
Dunhaga 3, IS-107 Reykjavik, Iceland

DESY 94-027
hep-th/9403027

Abstract

A repeatedly discussed gedanken experiment, proposed by Fermi to check Einstein causality, is reconsidered. It is shown that, contrary to a recent statement made by Hegerfeldt, there appears no causality paradoxon in a proper theoretical description of the experiment.

PACS number: 03.65.Bz

In a recent letter [1] Hegerfeldt discusses a gedanken experiment proposed by Fermi to determine the speed by which causal influences propagate. He argues that the theoretical description of this experiment in terms of transition probabilities leads to results which are in conflict with the existence of a maximal propagation speed c . Hegerfeldt suggests that the difficulties might disappear if one drops some implicit assumptions about the preparability of states with certain specific localization properties (points a) to c) of his conclusions). He does not settle the question whether the theory complies with Einstein causality, however.

In this letter we would like to set forth that there are no difficulties with Einstein causality in the theoretical setting of relativistic quantum field theory (RQFT). We will explain why, on one hand, transition probabilities are not a suitable tool for a thorough discussion of causal effects: what is required is a comparison of expectation values. On the other hand we will show that the points indicated by Hegerfeldt as possible loopholes to evade causality problems have to be taken seriously indeed and require a more careful analysis. Taking these facts into account we

will arrive at the conclusion that there is no conflict between the gedanken experiment of Fermi and the theoretical predictions of RQFT.

The experimental setup envisaged by Fermi to determine the propagation speed c of causal influences can be described as follows (cf. [2] and, for further references, [1]): one should prepare a state consisting of two atoms which are localized in disjoint regions separated by a distance R . One atom should be in its ground state, the other one in an excited state. If causal influences propagate with maximal velocity c one should not observe any impact of the excited atom on the atom in the ground state (e.g., by an emitted photon) within the time interval $0 < t < R/c$. Such events should be observed only at later times.

It is of importance that the atoms in this experiment have well defined localization properties. Hence in a theoretical discussion of the setup one first has to define in precise terms what one means by the statement that some physical state S (e.g., the state considered by Fermi, consisting of two atoms) looks at time $t = 0$, say, inside a region $\mathcal{R}$ like a given state G (e.g., like an atom in its ground state) [3]. From the point of view of physics the appropriate definition seems to be the following one: it is impossible to distinguish S from G by any measurement M which one performs at the given time in the region $\mathcal{R}$. In the theoretical setting this amounts to the requirement that the expectation values of all operators (observables) O_M corresponding to these measurements have to coincide. Hence if ψ_S, ψ_G denote the Hilbert space vectors representing S and G respectively there must hold

$$\langle \psi_S | O_M | \psi_S \rangle = \langle \psi_G | O_M | \psi_G \rangle. \quad (1)$$

This condition on ψ_S involves matrix elements of a multitude of observables and one may therefore ask whether it can be reformulated in terms of projection operators which test whether S coincides with G in the given region. The answer turns out to be different in relativistic and non-relativistic theories where the following alternatives hold.

i) Linear combinations of vectors ψ_S satisfying (1) again satisfy this condition (after normalization). This case is generic in non-relativistic quantum field theory, where it appears for a total set of states G . These states describe a situation where one has locally maximal information about the underlying system (locally pure states). Examples are the Fock vacuum and all coherent states. For any such state G one may consider the projection operator $O_{G \text{ inside } \mathcal{R} ?}$, projecting onto the subspace of the physical Hilbert space spanned by all vectors ψ_S satisfying condition (1). This projection operator can be used to decide whether some arbitrary state A coincides with G inside of $\mathcal{R}$: one simply has to calculate the transition probability $\langle \psi_A | O_{G \text{ inside } \mathcal{R} ?} | \psi_A \rangle$ and to check whether it is equal to 1. Thus in the non-relativistic setting there exist projection

operators which completely fix the local properties of states and it is then possible to study these properties in terms of transition probabilities.

ii) Certain normalized linear combinations of vectors ψ_S satisfying condition (1) do not comply with this condition. This is the case in RQFT for *every* choice of G . Phrased differently: all states G look locally like mixtures (of an, as a matter of fact, infinite number of states). In contrast to the non-relativistic case, it is thus not possible to fix the local properties of states with the help of projection operators [4].

In view of these facts one is forced to base the local analysis of states, which is fundamental in any discussion of causal effects, on a comparison of states in the sense of relation (1); transition probabilities are not the adequate tool to study this issue in relativistic theories. This important point may be illustrated by a simple example. If G is, e.g., the vacuum state in RQFT, then the expectation value $\langle \psi_G | O_M | \psi_G \rangle$ cannot vanish for any positive operator O_M corresponding to a localized measurement [5]. To test for a local deviation of S from G one can therefore not take as a criterion that the expectation value of some suitable projection operator (or, more generally, some positive operator) has a non-zero expectation value in the state S . For this expectation value would be non-zero even if the vacuum G is present. This point has been overlooked in [1] and led to the apparent paradoxa. A deviation of S from G would show up, however, in *different* values of the left and right sides of (1) for some O_M [6].

After these general remarks let us turn now to the actual discussion of Fermi's gedanken experiment. Let X be the ground state of an isolated atom which is localized in the vicinity of 0 and surrounded by vacuum and let ψ_X be the corresponding state vector. Following Fermi, we consider a state S , described by a vector ψ_S , which looks inside a ball $\mathcal{R}$ of radius R about 0 like X , i.e.,

$$\langle \psi_S | O_M | \psi_S \rangle = \langle \psi_X | O_M | \psi_X \rangle \quad (2)$$

for all observables O_M which are localized in $\mathcal{R}$. In the complement $\mathcal{R}^c$ of this ball S may look like any other state Y , e.g., like some excited atom. If, as expected, the subsystem in $\mathcal{R}^c$ does not affect the atom in $\mathcal{R}$ within the time interval $0 < t < R/c$ it should not be possible to discriminate S from X by any measurement M' which one performs at time t within the ball $\mathcal{R}_t$ of radius $R - ct$ about 0. Phrased differently: S should still look like an atom in its ground state within the smaller region $\mathcal{R}_t$. Hence, using the Heisenberg picture, there should hold in the theoretical setting

$$\langle \psi_S | O_{M'}(t) | \psi_S \rangle = \langle \psi_X | O_{M'}(t) | \psi_X \rangle, \quad (3)$$

where

$$O_{M'}(t) = e^{itH/\hbar} O_{M'} e^{-itH/\hbar}. \quad (4)$$

It is a fundamental fact that relation (3) is a consequence of relation (2) in theories where the underlying equations of motion are hyperbolic. Within the setting of RQFT this fact is called “primitive causality” [7] and has been established in models, cf. for example [8]. It is independent of the spectral properties of the generator H , which in fact depend on the systems which one considers (few body systems, thermal states, etc.). Hence in this respect the predictions of RQFT are in perfect agreement with the ideas of Fermi.

There remains, however, the question of whether the theory is capable of describing the physical situation envisaged by Fermi, cf. point c) in [1]. Given two vectors ψ_X , ψ_Y corresponding to states X , Y , does there exist a vector ψ_S describing the *composite* state S which looks like X in a given region $\mathcal{R}$ and like Y in its complement $\mathcal{R}^c$? These requirements fix ψ_S completely and can be cast into the following condition on the expectation values,

$$\langle \psi_S | O_M \cdot O_{M^c} | \psi_S \rangle = \langle \psi_X | O_M | \psi_X \rangle \langle \psi_Y | O_{M^c} | \psi_Y \rangle, \quad (5)$$

where O_M and O_{M^c} respectively denote operators corresponding to measurements in $\mathcal{R}$ and $\mathcal{R}^c$. Relation (5) gives formal expression to the idea that S is composed of states X and Y which are localized (in the sense of condition (1)) in disjoint regions and do not “overlap”, cf. point a) in [1].

The question of whether such product states exist is known in RQFT as the problem of “causal (statistical) independence” [9, 10]. It has an affirmative answer [11], but the vectors ψ_S have in general infinite energy even if ψ_X and ψ_Y have finite energy. This phenomenon can be traced back to the uncertainty principle and may be easily understood in the framework of non-relativistic quantum mechanics: if ψ_X and ψ_Y are the configuration space wave functions of distinguishable systems then the wave function ψ_S of the composite state is given by

$$\psi_S(\mathbf{x}, \mathbf{y}) = N \cdot \begin{cases} \psi_X(\mathbf{x}) \psi_Y(\mathbf{y}) & \text{for } \mathbf{x} \in \mathcal{R}, \mathbf{y} \in \mathcal{R}^c \\ 0 & \text{otherwise} \end{cases} \quad (6)$$

where N is a normalization constant. This function has in general a discontinuity when $\mathbf{x}$ or $\mathbf{y}$ are at the boundary of $\mathcal{R}$, unless the wave functions ψ_X and ψ_Y happen to vanish at these points. (Note that wave functions of states with sharp energy, such as bound states, in general do not have such nodes.) As a consequence, the expectation value of the Hamiltonian becomes infinite. In RQFT the situation is even worse because of pair creation. There it turns out

that due to such processes the vector ψ_S cannot be an element of the physical Hilbert space $\mathcal{H}$ describing few body systems (e.g., Fock space in free field theory).

Thus the theory predicts that every member of the ensemble described by S has infinite energy. Hence a preparation of this state would not be possible in practice. This fact seems to be in conflict with the ideas of Fermi, but the apparent difficulty disappears if one notices that for the determination of c it suffices to consider “tame” states T which look like X in a region $\mathcal{R}_<$ and like Y in $\mathcal{R}_>$, where $\mathcal{R}_<$ is slightly smaller and $\mathcal{R}_>$ slightly larger than $\mathcal{R}$. It is not really necessary to completely fix the state T in the layer between these two regions. By making this layer sufficiently small one can then determine c , as outlined above, with arbitrary precision.

It has been shown in RQFT under very general conditions that there exist vectors ψ_T in the physical Hilbert space $\mathcal{H}$ which satisfy condition (5) for the slightly smaller regions [12, 13]. The existence of such vectors has also been established in models [14, 15, 16]. Thus also in this respect Fermi’s gedanken experiment poses no theoretical problems.

It should be mentioned that the vectors ψ_T have to be carefully adjusted in the layer between the two regions $\mathcal{R}_<$ and $\mathcal{R}_>$ in order to become elements of the physical Hilbert space $\mathcal{H}$. This fact may be viewed as the process of “renormalization”, indicated in point b) of [1], which surrounds state X by some “cloud”. In more physical terms: any state of the type considered by Fermi which can actually be prepared in an experiment necessarily contains, besides the two atoms, other particles, e.g., photons. This fact is the basic reason for the apparently non-causal behaviour of transition probabilities, discussed in [1]. But it is not in conflict with the existence of a maximal propagation speed c .

References

- [1] G.C. Hegerfeldt, Phys. Rev. Lett. **72** (1994) 596
- [2] E. Fermi, Rev. Mod. Phys. **4** (1932) 87
- [3] We assume that G exists as a global state described by a vector in the physical Hilbert space. This assumption causes no problems if one considers configurations of atoms. But it would require further discussions if one thinks, e.g., of quarks as subsystems of hadrons (confinement).
- [4] These marked differences between relativistic and non-relativistic theories are actually encoded in the specific structure of the algebras generated by the respective observables in bounded regions. In non-relativistic quantum field theory these algebras are generically of

type I according to the classification scheme of von Neumann. In RQFT they are of *type III*, cf. for example

R. Haag, *Local Quantum Physics*, (Springer, Berlin Heidelberg 1992)

- [5] This is the statement of the Reeh-Schlieder-Theorem, cf. the preceding reference. As a matter of fact the statement holds for almost any physical state G (the set of such states is of *second category*).
- [6] The comparison of matrix elements of observables seems, in a discussion of causal effects, also natural from a more general point of view: if one were to determine the effect of a perturbation of a state ϕ with the help of some observable Q , one would *compare* the moments of Q in the unperturbed state ϕ and the perturbed state ϕ' , say. The differences $D_n = \langle \phi' | Q^n | \phi' \rangle - \langle \phi | Q^n | \phi \rangle$, $n \in \mathbf{N}$ would then provide a measure for the observed effect of the perturbation.
- [7] R. Haag, B. Schroer, J. Math. Phys. **3** (1962) 248
- [8] J. Glimm, A. Jaffe, in *Statistical Mechanics and Quantum Field Theory*, Les Houches 1970, (Gordon and Breach, New York 1971), p. 1
- [9] R. Haag, D. Kastler, J. Math. Phys. **5** (1964) 848
- [10] S. Schlieder, Commun. Math. Phys. **13** (1969) 216
- [11] H. Roos, Commun. Math. Phys. **16** (1970) 238
- [12] D. Buchholz, E.H. Wichmann, Commun. Math. Phys. **106** (1986) 321
- [13] D. Buchholz, C. D'Antoni, K. Fredenhagen, Commun. Math. Phys. **111** (1987) 123
- [14] D. Buchholz, Commun. Math. Phys. **36** (1974) 287
- [15] W. Driessler, Commun. Math. Phys. **70** (1979) 213
- [16] S. Summers, Commun. Math. Phys. **86** (1982) 111